


HOMFLYPT homology for links in handlebodies


Or: All I know about Artin–Tits groups; and a filler for the remaining 59 minutes


Daniel Tubbenhauer





Joint with David Rose


January 2019









































































1 Links and braids in handlebodies

- Braid diagrams
- Links in handlebodies

2 Some “low-genus-coincidences”


- The ball and the torus
- The torus and the double torus

3 Arbitrary genus


- Braid invariants – some ideas
- Link invariants – some ideas

Let $\text{Br}(g, n)$ be the group defined as follows.


Generators. Braid and twist generators



Relations. ► Reidemeister braid relations, type C relations and special relations, e.g.




Involves three players and inverses!



Let $\text{Br}(g, n)$ be the group defined as follows.

Generators. Braid and twist

$\ell_i \leftrightarrow$




Relations.

► Reidemeister braid relation


and special relations, e.g.

Involves three players and inverses!



$$\ell_1 t_2 \ell_1 t_2 = t_2 \ell_1 t_2 \ell_1$$

&





$$(\ell_1 t_2 \ell_1^{-1}) t_3 = t_3 (\ell_1 t_2 \ell_1^{-1})$$

Let $\text{Br}(g, n)$ be the group defined as follows.

Generators. Braid and twist

$\sigma_i \leftrightarrow$



Relations.


► Reidemeister braid relation

and special relations, e.g.

Involves three players and inverses!



=



=



$$\sigma_1 t_2 \sigma_1 t_2 = t_2 \sigma_1 t_2 \sigma_1$$

$$(\sigma_1 t_2 \sigma_1^{-1}) t_3 = t_3 (\sigma_1 t_2 \sigma_1^{-1})$$

Let $\text{Br}(g, n)$ be the group defined as follows.

Generators. Braid and twist generators

1 g 1 i i+1 n 1 i g 1 2 n

Fact (type A embedding).

$\text{Br}(g, n)$ is a subgroup of the usual braid group $\mathcal{B}\text{r}(g+n)$.


Relatio

A visualization exercise.

$\ell_1 t_2 \ell_1^{-1} t_2 = t_2 \ell_1 t_2 \ell_1^{-1}$ $(\ell_1 t_2 \ell_1^{-1}) t_3 = t_3 (\ell_1 t_2 \ell_1^{-1})$


The group $\mathcal{B}r(g, n)$ of braid in a g -times punctures disk $\mathcal{D}_g^2 \times [0, 1]$:

Two types of braidings, the usual ones and “winding around cores”, e.g.



Theorem (Häring-Oldenburg–Lambropoulou ~2002, Vershinin ~1998).


The map



is an isomorphism of groups $\text{Br}(g, n) \rightarrow \mathcal{B}\text{r}(g, n)$.

The group $\mathcal{B}r(g, n)$ of braid in a g -times punctures disk $\mathcal{D}_g^2 \times [0, 1]$:

Two types of embeddings, e.g.




The group $\mathcal{B}r(g, n)$ of braid in a g -times punctures disk $\mathcal{D}_g^2 \times [0, 1]$:

Two types of braidings, the usual ones and “winding around cores” e.g.

Note.


For the proof it is crucial that \mathcal{D}_g^2 and the boundary points of the braids \bullet are only defined up to isotopy, e.g.




\Rightarrow one can always “conjugate cores to the left”.

This is useful to define $\mathcal{B}r(g, \infty)$.

The Alexander closure on $\mathcal{B}r(g, \infty)$ is given by merging core strands at infinity.



wrong closure




correct closure

This is different from the [classical](#) Alexander closure.

The Alexander closure on $\mathcal{B}r(g, \infty)$ is given by merging core strands at infinity.

Theorem (Lambropoulou ~1993).

For any link ℓ in the genus g handlebody \mathcal{H}_g there is a braid in $\mathcal{B}r(g, \infty)$ whose (correct!) closure is isotopic to ℓ .



This is different from the [classical](#) Alexander closure.


The Alexander closure on $\mathcal{B}r(g, \infty)$ is given by merging core strands at infinity.

Theorem (Lambropoulou ~1993).


For any link ℓ in the genus g handlebody \mathcal{H}_g there is a braid in $\mathcal{B}r(g, \infty)$ whose (correct!) closure is isotopic to ℓ .

Fact.


\mathcal{H}_g is given by a complement in the 3-sphere \mathcal{S}^3 by an open tubular neighborhood of the embedded graph obtained by gluing $g + 1$ unknotted "core" edges to two vertices.



the 3-ball $\mathcal{H}_0 = \mathcal{D}^3$



a torus \mathcal{H}_1



\mathcal{H}_2


This is

The Markov moves on $\mathcal{B}r(g, \infty)$ are conjugation and stabilization.

Conjugation.

$$\ell \sim s\ell s^{-1}$$

for $\ell \in \mathcal{B}r(g, n)$, $s \in \langle \ell_1, \dots, \ell_{n-1} \rangle$




Stabilization.

$$(c\uparrow)\ell_n(\ell\uparrow)$$

$$\sim c\ell \sim (c\uparrow)\ell_n^{-1}(\ell\uparrow)$$

for $\ell, c \in \mathcal{B}r(g, n)$,




They are weaker than the [classical](#) Markov moves.

Theorem (Häring-Oldenburg–Lambropoulou ~2002).

Two links in \mathcal{H}_g are equivalent if and only if they are equal in $\mathcal{B}r(g, \infty)$ up to conjugation and stabilization.

$$\ell \sim s\ell s^{-1}$$


for $\ell \in \mathcal{B}r(g, n)$, $s \in \langle \ell_1, \dots, \ell_{n-1} \rangle$



Stabilization.

$$(c\uparrow)\ell_n(\ell\uparrow) \sim c\ell \sim (c\uparrow)\ell_n^{-1}(\ell\uparrow)$$

for $\ell, c \in \mathcal{B}r(g, n)$,





They are weaker than the classical Markov moves.

The Markov moves on $\mathcal{B}r(g, \infty)$ are conjugation and stabilization

Theorem (Häring-Oldenburg–Lambropoulou ~2002).

Two links in \mathcal{H}_g are equivalent if and only if they are equal in $\mathcal{B}r(g, \infty)$ up to conjugation and stabilization.

Example.



for $\beta \in \mathcal{B}r$

Stabilization

$(c\uparrow)\beta$

$\sim c\beta \sim (c\downarrow)\beta$


for $\beta, c \in \mathcal{B}r$

They are weakly

The Markov moves on $\mathcal{B}r(g, \infty)$ are conjugation and stabilization.


Conjugation.

$\ell \sim s\ell s^{-1}$
for $\ell \in \mathcal{B}r(g, n)$,



Stabilization.

$(c\uparrow)\ell_n(\ell\uparrow)$
 $\sim c\ell \sim (c\uparrow)\ell_n^{-1}(\ell\uparrow)$
for $\ell, c \in \mathcal{B}r(g, n)$,




They are weaker than the classical Markov moves.

The Markov moves on $\mathcal{B}r(g, \infty)$ are conjugation and stabilization.

Conjugation.


$\ell \sim s\ell s^{-1}$
for $\ell \in \mathcal{B}r(g, n)$,



Stabilization.

$(c\uparrow)\ell_n(\ell\uparrow)$
 $\sim c\ell \sim (c\uparrow)\ell_n^{-1}(\ell\uparrow)$
for $\ell, c \in \mathcal{B}r(g, n)$,

Let me explain what we can do.



They are weaker than the classical Markov moves.

Let Γ be a Coxeter graph.

Artin \sim 1925, **Tits** \sim 1961++. The Artin–Tits group and its Coxeter group quotient are given by generators–relations:

$$\begin{aligned} \text{AT}(\Gamma) &= \langle \ell_i \mid \underbrace{\cdots \ell_i \ell_j \ell_i}_{m_{ij} \text{ factors}} = \underbrace{\cdots \ell_j \ell_i \ell_j}_{m_{ij} \text{ factors}} \rangle \\ &\downarrow \\ \text{W}(\Gamma) &= \langle \sigma_i \mid \sigma_i^2 = 1, \underbrace{\cdots \sigma_i \sigma_j \sigma_i}_{m_{ij} \text{ factors}} = \underbrace{\cdots \sigma_j \sigma_i \sigma_j}_{m_{ij} \text{ factors}} \rangle \end{aligned}$$

Artin–Tits groups [▶ generalize](#) classical braid groups, Coxeter groups [▶ generalize](#) polyhedron groups.

$\cos(\pi/3)$ on a line:

type A_{n-1} : $1 \text{ --- } 2 \text{ --- } \dots \text{ --- } n-2 \text{ --- } n-1$

The classical case. Consider the map

$$\beta_i \mapsto \begin{array}{cccc} 1 & i & i+1 & n \\ \uparrow & \nearrow & \nearrow & \uparrow \\ \dots & \times & \dots & \\ \uparrow & \searrow & \searrow & \uparrow \\ 1 & i & i+1 & n \end{array} \quad \text{braid rel.:} \quad \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \\ \diagdown \quad \diagup \quad \diagdown \\ \diagup \quad \diagdown \quad \diagup \\ \uparrow \quad \uparrow \quad \uparrow \end{array} = \begin{array}{c} \uparrow \quad \uparrow \quad \uparrow \\ \diagup \quad \diagdown \quad \diagup \\ \diagdown \quad \diagup \quad \diagdown \\ \uparrow \quad \uparrow \quad \uparrow \end{array}$$

Artin ~1925. This gives an isomorphism of groups $AT(A_{n-1}) \xrightarrow{\cong} \mathcal{B}r(0, n)$.

$\cos(\pi/3)$ on a line:

Jones ~1987.

Markov trace on the Hecke algebra of type A

\rightsquigarrow two variable q, a polynomial invariant (HOMFLYPT polynomial).

The clas

q =Hecke parameter ; a =trace parameter .



braid rel.:



Artin ~1925. This gives an isomorphism of groups $AT(A_{n-1}) \xrightarrow{\cong} \mathcal{B}r(0, n)$.

I will come back to this with more details for general genus g .
For the time being: This works quite well!

$\cos(\pi/3)$ on a line:

Jones ~1987.

Markov trace on the Hecke algebra of type A

↪ two variable q, a polynomial invariant (HOMFLYPT polynomial).

The clas

q =Hecke parameter ; a =trace parameter .

Khovanov ~2005; categorification.

Hochschild homology on complexes of the Hecke category of type A

↪ “three variable q, t, a homological invariant” (HOMFLYPT homology).

q =Hecke parameter ; t =homological parameter ; a =Hochschild parameter .


Artin ~1929. This gives an isomorphism of groups $AI(A_{n-1}) \cong \mathcal{B}I(0, n)$.

I will come back to this with more details for general genus g .
For the time being: This works quite well!

$\cos(\pi/3)$ on a circle.



Affine adds genus. Consider the map



tom Dieck ~1998. (Earlier reference?) This gives an isomorphism of groups $\mathbb{Z} \times \text{AT}(\tilde{A}_{n-1}) \xrightarrow{\cong} \mathcal{B}r(1, n)$.

$\cos(\pi/3)$ on a circle.


Orellana–Ram ~2004. (Earlier reference?)

Markov trace on the Hecke algebra of type \tilde{A}

Affine a

\rightsquigarrow two variable q, a polynomial invariant (HOMFLYPT polynomial).

q =Hecke parameter ; a =trace parameter .



tom Dieck ~1998. (Earlier reference?) This gives an isomorphism of groups

$\mathbb{Z} \times \text{AT}(\tilde{A}_{n-1}) \cong \mathcal{Rr}(1, n)$

I will come back to this with more details for general genus g .
For the time being: This works quite well!

$\cos(\pi/3)$ on a circle.

Orellana–Ram ~2004. (Earlier reference?)

Markov trace on the Hecke algebra of type \tilde{A}

↪ two variable q, a polynomial invariant (HOMFLYPT polynomial).

q =Hecke parameter ; a =trace parameter .

Affine a

???; categorification.

Hochschild homology on complexes of the Hecke category of type \tilde{A}

↪ “three variable q, t, a homological invariant” (HOMFLYPT homology).

q =Hecke parameter ; t =homological parameter ; a =Hochschild parameter .

tom Dieck ~1996. (Earlier reference?) This gives an isomorphism of groups

$\mathbb{Z} \times \text{AT}(\tilde{A}_{n-1}) \cong \text{Br}(1, n)$

I will come back to this with more details for general genus g .
For the time being: This works quite well!


$\cos(\pi/3)$ on a circle.

Fact. One can recover the (missing) generator of \mathbb{Z} if one works with extended affine type A.


Affine adds

$\beta_0 \mapsto$



"extended, extra generator" \mapsto



and



give



tom Dieck


$$\mathbb{Z} \times \text{AT}(\tilde{A}_{n-1}) \xrightarrow{\cong} \mathcal{B}r(1, n).$$

m of groups

$\cos(\pi/4)$ on a line:

$$\text{type } C_n: 0 \stackrel{4}{=} 1 - 2 - \dots - n-1 - n$$

The semi-classical case. Consider the map



Brieskorn \sim 1973. This gives an isomorphism of groups $\text{AT}(C_n) \xrightarrow{\cong} \mathcal{B}r(1, n)$.


$\cos(\pi/4)$ on a line:

Geck–Lambropoulou ~1997.

Markov trace on the Hecke algebra of type C

\rightsquigarrow two variable q, a polynomial invariant (HOMFLYPT polynomial).

q =Hecke parameter ; a =trace parameter .



Brieskorn ~1973. This gives an isomorphism of groups $AT(C_n) \xrightarrow{\cong} \mathcal{B}r(1, n)$.

I will come back to this with more details for general genus g .
For the time being: This works quite well!

$\cos(\pi/4)$ on a line:

Geck–Lambropoulou ~1997.

Markov trace on the Hecke algebra of type C

\rightsquigarrow two variable q , a polynomial invariant (HOMFLYPT polynomial).

q =Hecke parameter ; a =trace parameter .

1 2 n 1 i i+1 n

Rouquier ~2012, Webster–Williamson ~2009; categorification.


Hochschild homology on complexes of the Hecke category of type C

\rightsquigarrow “three variable q , t , a homological invariant” (HOMFLYPT homology).

q =Hecke parameter ; t =homological parameter ; a =Hochschild parameter .

Brieskorn ~1975. This gives an isomorphism of groups $A1(\mathbb{C}_n) \rightarrow \mathcal{S}1(1, n)$.

I will come back to this with more details for general genus g .
For the time being: This works quite well!



$\cos(\pi/4)$ on a line:

Fact. (Not true in type A.)

There is a whole infinite family of Markov traces,
one for each choice of a value for essential unlinks.

The

β_0

extra parameter and extra parameter etc.

However, I only know the categorification of one of these.

Brieskorn \sim 1973. This gives an isomorphism of groups $AT(C_n) \xrightarrow{\cong} \mathcal{B}r(1, n)$.

$\cos(\pi/4)$ on a line:

Fact. (Not true in type A.)

There is a whole infinite family of Markov traces, one for each choice of a value for essential unlinks.

The β_0

extra parameter and extra parameter etc.

However, I only know the categorification of one of these.

Fact. (Not true in type A.)

Brieskorn ~ 1973 . There is also a second Hecke parameter, $\mathbb{C}[n] \xrightarrow{\mathbb{R}} \mathcal{B}r(1, n)$, which we do not know how to categorify yet.

$\cos(\pi/4)$ twice on a line:

$$\text{type } \tilde{C}_n: 0^1 \underline{\underline{4}} 1 \text{ --- } 2 \text{ --- } \dots \text{ --- } n-1 \text{ --- } n \underline{\underline{4}} 0^2$$


Affine adds genus. Consider the map

$$\beta_{0^1} \mapsto \begin{array}{c} \color{red}{1} \quad \color{red}{1} \quad \color{red}{n} \quad \color{red}{2} \\ \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \\ \color{red}{1} \quad \color{red}{1} \quad \color{red}{n} \quad \color{red}{2} \end{array} \quad \& \quad \beta_i \mapsto \begin{array}{c} i \quad i+1 \\ \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \\ i \quad i+1 \end{array} \quad \& \quad \beta_{0^2} \mapsto \begin{array}{c} \color{red}{1} \quad \color{red}{1} \quad \color{red}{n} \quad \color{red}{2} \\ \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \\ \color{red}{1} \quad \color{red}{1} \quad \color{red}{n} \quad \color{red}{2} \end{array}$$


Allcock ~1999. This gives an isomorphism of groups $\text{AT}(\tilde{C}_n) \xrightarrow{\cong} \mathcal{B}r(2, n)$.

This case is strange – it only arises under conjugation:


$\cos(\pi/4)$ twice



By a miracle, one can avoid the special relation



This relation involves three players and inverses. Bad!




$\beta_{01} \mapsto$


Allcock \sim 1999. This gives an isomorphism of groups $\text{AT}(\tilde{C}_n) \xrightarrow{\cong} \mathcal{B}r(2, n)$.

This case is strange – it only arises under conjugation:


$\cos(\pi/4)$ twice



By a miracle, one can avoid the special relation



This relation involves three players and inverses. Bad!




$\beta_{01} \mapsto$

Currently, not much seems to be known, but I think the same story works.


Allcock \sim 1999. This gives an isomorphism of groups $\text{AT}(\tilde{C}_n) \xrightarrow{\cong} \mathcal{B}r(2, n)$.

This case is strange – it only arises under conjugation:


$\cos(\pi/4)$ twice



By a miracle, one can avoid the special relation



This relation involves three players and inverses. Bad!




$\beta_{01} \mapsto$

Affine adds ge

Currently, not much seems to be known, but I think the same story works.

Allcock

However, this is where it seems to end, e.g. genus $g = 3$ wants to be n).



But the special relation makes it a mere quotient.
So: In the remaining time I tell you what works.

$\cos(\pi/4)$ twice on a line:

Currently known (to the best of my knowledge).


Genus	type A	type C
$g = 0$	$\mathcal{B}r(n) \cong AT(A_{n-1})$	
$g = 1$	$\mathcal{B}r(1, n) \cong \mathbb{Z} \times AT(\tilde{A}_{n-1}) \cong AT(\hat{A}_{n-1})$	$\mathcal{B}r(1, n) \cong AT(C_n)$
$g = 2$		$\mathcal{B}r(2, n) \cong AT(\tilde{C}_n)$
$g \geq 3$		

And some $\mathbb{Z}/2\mathbb{Z}$ -orbifolds ($\mathbb{Z}/\infty\mathbb{Z}$ = puncture):


Genus	type D	type B
$g=0$		
$g=1$	$\mathcal{B}r(1, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \mathbb{Z}/2\mathbb{Z} \times AT(D_n)$	$\mathcal{B}r(1, n)_{\mathbb{Z}/\infty\mathbb{Z}} \cong AT(B_n)$
$g=2$	$\mathcal{B}r(2, n)_{\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}} \cong (\mathbb{Z}/2\mathbb{Z})^2 \times AT(\tilde{D}_n)$	$\mathcal{B}r(2, n)_{\mathbb{Z}/\infty\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}} \cong \mathbb{Z}/2\mathbb{Z} \times AT(\tilde{B}_n)$
$g \geq 3$		

(For orbifolds "genus" is just an analogy.)


Philosophy 1: Reshetikhin–Turaev with “huge” colors.




Philosophy 1: Reshetikhin–Turaev with “huge” colors.




Philosophy 1: Reshetikhin–Turaev with “huge” colors.




Philosophy 1: Reshetikhin–Turaev with “huge” colors.




Philosophy 1: Reshetikhin–Turaev with “huge” colors.




Philosophy 1: Reshetikhin–Turaev with “huge” colors.




Philosophy 1: Reshetikhin–Turaev with “huge” colors.



Philosophy 1: Reshetikhin–Turaev with “huge” colors.




Philosophy 1: Reshetikhin–Turaev with “huge” colors.




Philosophy 1: Resh

Note that the type A embedding guarantees that any usual invariant of braids produces an invariant of braids in \mathcal{H}_g .




Note that the type A embedding guarantees that any usual invariant of braids produces an invariant of braids in \mathcal{H}_g .



Genus $g = 0, 1$.

Works quite well (e.g. use Naisse-Vaz's ideas on the categorified level).

Note that the type A embedding guarantees that any usual invariant of braids produces an invariant of braids in \mathcal{H}_g .



Genus $g = 0, 1$.

Works quite well (e.g. use Naisse-Vaz's ideas on the categorified level).

We mimic this for M being "huge, but finite".

Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$.

Tuples $\mathbf{I} = (k_1, \dots, k_N) \in \mathbb{N}_{\geq 1}^N$ with $k_1 + \dots + k_N = N \iff$ parabolic subgroups

$$W_{\mathbf{I}} = W(A_{k_1-1}) \times \dots \times W(A_{k_N-1}) \subset W.$$

W acts on $\mathbb{R} = \mathbb{R}_N = \mathbb{k}[x_1, \dots, x_N]$ via permutation \rightsquigarrow rings of invariants $\mathbb{R}^{\mathbf{I}}$.

Bimodules. Identities, restriction (“merge”) and induction (“split”), e.g.

$$\begin{array}{c} 1 & 1 & 1 \\ | & | & | \\ 1 & 1 & 1 \end{array} \iff \mathbb{R}^{(1,1,1)} = \mathbb{R}, \quad \begin{array}{c} 2 & 1 \\ | & | \\ 2 & 1 \end{array} \iff \mathbb{R}^{(2,1)} = \mathbb{R}^{\sigma_1} = \mathbb{k}[x_1 + x_2, x_1 x_2, x_3].$$

$$\begin{array}{c} k+l \\ \cup \\ k \quad l \end{array} \iff \text{shift} \mathbb{R}^{(k+l)} \otimes_{\mathbb{R}^{(k+l)}} \mathbb{R}^{(k,l)}, \quad \begin{array}{c} k \quad l \\ \cup \\ k+l \end{array} \iff \mathbb{R}^{(k,l)} \otimes_{\mathbb{R}^{(k+l)}} \mathbb{R}^{(k+l)}.$$

Define $\mathcal{S}_s^q(W)$ as the full 2-subcategory of the rings&bimodules 2-category.

Singular Soergel bimodules $\mathcal{S}_s^{\mathfrak{q}}(W)$ for $W = W(A_{N-1})$.

Tuples $\mathbf{I} = (k_1, \dots, k_N) \in \mathbb{N}_{\geq 1}^N$ with $k_1 + \dots + k_N = N \iff$ parabolic subgroups

$$W_{\mathbf{I}} = W(A_{k_1-1}) \times \dots \times W(A_{k_N-1}) \subset W.$$

W acts on $\mathbb{R} = \mathbb{R}_N = \mathbb{k}[x_1, \dots, x_N]$. Rings of invariants $\mathbb{R}^{\mathbf{I}}$.

Everything is \mathbb{Z} -graded, called \mathfrak{q} -grading.
I just omit this for simplicity.

Bimodules. Identities, restriction (“merge”) and induction (“split”), e.g.

$$\begin{array}{c} 1 & 1 & 1 \\ | & | & | \\ 1 & 1 & 1 \end{array} \iff \mathbb{R}^{(1,1,1)} = \mathbb{R}, \quad \begin{array}{c} 2 & 1 \\ | & | \\ 2 & 1 \end{array} \iff \mathbb{R}^{(2,1)} = \mathbb{R}^{\sigma_1} = \mathbb{k}[x_1 + x_2, x_1 x_2, x_3].$$

$$\begin{array}{c} k+l \\ \cup \\ k \quad l \end{array} \iff \text{shift} \mathbb{R}^{(k+l)} \otimes_{\mathbb{R}^{(k+l)}} \mathbb{R}^{(k,l)}, \quad \begin{array}{c} k \quad l \\ \cup \\ k+l \end{array} \iff \mathbb{R}^{(k,l)} \otimes_{\mathbb{R}^{(k+l)}} \mathbb{R}^{(k+l)}.$$

Define $\mathcal{S}_s^{\mathfrak{q}}(W)$ as the full 2-subcategory of the rings&bimodules 2-category.

Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$.

A monoidal structure is given by

$$\begin{array}{c} 1 & & 1 \\ & \cup & \\ & & \\ & \cap & \\ & & \\ 1 & & 1 \end{array} = \begin{array}{c} 2 \\ | \\ 1 & & 1 \end{array} \leftarrow \text{glue} \rightarrow \begin{array}{c} 1 & & 1 \\ & \cup & \\ & & \\ & \cap & \\ & & \\ 1 & & 2 \end{array} \iff R \otimes_{R^{\sigma_1}} R \cong R \otimes_{R^{\sigma_1}} R^{\sigma_1} \otimes_{R^{\sigma_1}} R.$$

This gives a way to define bimodules associated to any web built out of merge and split.

Bimodules. Identities, restriction (“merge”) and induction (“split”), e.g.

$$\begin{array}{c} 1 & 1 & 1 \\ | & | & | \\ 1 & 1 & 1 \end{array} \iff R^{(1,1,1)} = R, \quad \begin{array}{c} 2 & 1 \\ | & | \\ 2 & 1 \end{array} \iff R^{(2,1)} = R^{\sigma_1} = \mathbb{k}[x_1 + x_2, x_1 x_2, x_3].$$

$$\begin{array}{c} k+l \\ | \\ k & & l \end{array} \iff \text{shift} R^{(k+l)} \otimes_{R^{(k+l)}} R^{(k,l)}, \quad \begin{array}{c} k & l \\ & \cup \\ & & \\ & \cap \\ & & \\ k+l \end{array} \iff R^{(k,l)} \otimes_{R^{(k+l)}} R^{(k+l)}.$$

Define $\mathcal{S}_s^q(W)$ as the full 2-subcategory of the rings&bimodules 2-category.

Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$.

A monoidal structure is given by

$$\begin{array}{c} 1 & 1 \\ \diagdown & / \\ & \text{---} \\ / & \diagdown \\ 1 & 1 \end{array} = \begin{array}{c} 2 \\ | \\ 1 & 1 \end{array} \leftarrow \text{glue} \rightarrow \begin{array}{c} 1 & 1 \\ / & \diagdown \\ & \text{---} \\ | \\ 2 \end{array} \iff R \otimes_{R^{\sigma_1}} R \cong R \otimes_{R^{\sigma_1}} R^{\sigma_1} \otimes_{R^{\sigma_1}} R.$$

This gives a way to define bimodules associated to any web built out of merge and split.

Bimodules. Identify (There are several bimodule isomorphisms, e.g. "split"), e.g.

$$\begin{array}{c} 1 & 1 & 1 \\ | & | & | \\ 1 & 1 & 1 \end{array} \iff R \otimes R \otimes R$$

$$\begin{array}{c} k+l+m \\ | \\ \diagdown & / \\ & \text{---} \\ / & \diagdown \\ k & l & m \end{array} \cong \begin{array}{c} k+l+m \\ | \\ \diagdown & / \\ & \text{---} \\ / & \diagdown \\ k & l & m \end{array} \quad \& \quad \begin{array}{c} k & l & m \\ / & \diagdown \\ & \text{---} \\ | \\ k+l+m \end{array} \cong \begin{array}{c} k & l & m \\ / & \diagdown \\ & \text{---} \\ | \\ k+l+m \end{array}$$

$$\begin{array}{c} k+l \\ | \\ \diagdown & / \\ & \text{---} \\ / & \diagdown \\ k & l \end{array} \iff \text{shift}$$

Hence, we can unambiguously write

$$\begin{array}{c} k_1 + \dots + k_r \\ | \\ \diagdown & / \\ & \text{---} \\ / & \diagdown \\ k_1 & \dots & k_r \end{array} \quad \& \quad \begin{array}{c} k_1 & k_r \\ / & \diagdown \\ & \text{---} \\ | \\ k_1 + \dots + k_r \end{array}$$

Define $\mathcal{S}_s^q(W)$ as which one could call thick merge and split. $\mathcal{S}_s^q(W)$ is a 2-category.

Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$.

Soergel ~1992, Williamson ~2010.

Tuples $I = (i_1, \dots, i_N) \in \{1, \dots, N\}^N$ subgroups

$$W_I = W(A_{k_1-1}) \times \cdots \times W(A_{k_N-1}) \subset W.$$

W acts on $R = R_N = \mathbb{k}[x_1, \dots, x_N]$ via permutation \rightsquigarrow rings of invariants R^I .

Bimodules. Identities, restriction (“merge”) and induction (“split”), e.g.

$$\begin{array}{c} 1 & 1 & 1 \\ | & | & | \\ 1 & 1 & 1 \end{array} \iff R^{(1,1,1)} = R, \quad \begin{array}{c} 2 & 1 \\ | & | \\ 2 & 1 \end{array} \iff R^{(2,1)} = R^{\sigma_1} = \mathbb{k}[x_1 + x_2, x_1 x_2, x_3].$$

$$\begin{array}{c} k+l \\ | \\ \text{---} \\ | & | \\ k & l \end{array} \iff \text{shift} R^{(k+l)} \otimes_{R^{(k+l)}} R^{(k,l)}, \quad \begin{array}{c} k & l \\ \text{---} \\ | \\ k+l \end{array} \iff R^{(k,l)} \otimes_{R^{(k+l)}} R^{(k+l)}.$$

Define $\mathcal{S}_s^q(W)$ as the full 2-subcategory of the rings&bimodules 2-category.

Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$.

Soergel ~1992, Williamson ~2010.

Tuples $\Gamma = (\gamma_1, \dots, \gamma_n) \in \{1, 2\}^n$ categorifies the Hecke algebra (or rather, the algebroid) of subgroups

Rouquier ~2004, Mackaay–Stošić–Vaz ~2008, Webster–Williamson ~2009, etc.

There are certain complex (“t-graded”) of singular Soergel bimodules, e.g.

$$[\beta_i]_M = \begin{array}{c} l \quad k \\ \diagdown \quad \diagup \\ k \quad l \end{array} = \begin{array}{c} k-l \\ \diagdown \quad \diagup \\ k \quad l \\ 0 \end{array} \xrightarrow{d_0^+} \mathbf{qt} \begin{array}{c} k-l \\ \diagdown \quad \diagup \\ k \quad l \\ 1 \end{array} \xrightarrow{d_1^+} \dots \xrightarrow{d_{l-1}^+} \mathbf{q}^l \mathbf{t}^l \begin{array}{c} k \\ \diagdown \quad \diagup \\ k \quad l \end{array}$$

providing a categorical action of the Artin–Tits group of type A.

1 1 1

2 1



$$\iff \text{shift} R^{(k+l)} \otimes_{R^{(k+l)}} R^{(k,l)},$$



$$\iff R^{(k,l)} \otimes_{R^{(k+l)}} R^{(k+l)}.$$

Define $\mathcal{S}_s^q(W)$ as the full 2-subcategory of the rings&bimodules 2-category.

Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$.

Soergel ~1992, Williamson ~2010.

Tuples $\mathbf{I} = (i_1, \dots, i_n) \in \{1, \dots, N\}^n$ categorifies the Hecke algebra (or rather, the algebroid) of subgroups

Rouquier ~2004, Mackaay–Stošić–Vaz ~2008, Webster–Williamson ~2009, etc.

There are certain complex (“t-graded”) of singular Soergel bimodules, e.g.

$$[\beta_i]_M = \begin{array}{c} l \quad k \\ \diagdown \quad \diagup \\ k \quad l \end{array} = \begin{array}{c} k-l \\ \diagdown \quad \diagup \\ k \quad l \\ 0 \end{array} \xrightarrow{d_0^+} \mathbf{qt} \begin{array}{c} k-l \\ \diagdown \quad \diagup \\ k \quad l \\ +1 \\ 1 \end{array} \xrightarrow{d_1^+} \dots \xrightarrow{d_{l-1}^+} \mathbf{q}^l \mathbf{t}^l \begin{array}{c} k \\ \diagdown \quad \diagup \\ k \quad l \end{array}$$

providing a categorical action of the Artin–Tits group of type A.

1 1 1 2 1

Hence, we are in business by taking $M \gg n$:

$$\uparrow \iff \begin{array}{c} 1 \\ | \\ 1 \end{array} \quad \text{and} \quad \begin{array}{c} | \\ | \\ | \end{array} \iff \begin{array}{c} M \\ | \\ M \end{array} = \begin{array}{c} M \\ | \\ M \end{array} \quad \text{and} \quad \begin{array}{c} \uparrow \\ | \\ \downarrow \end{array} \iff \begin{array}{c} M \quad 1 \\ \diagdown \quad \diagup \\ M \quad 1 \end{array} \quad \text{etc.}$$

Defin This defines a fairly strong complex-valued invariant $[\mathcal{L}]_M$ of $\mathcal{L} \in \mathcal{B}r(g, n)$.

Partial Hochschild homology (à la Hogancamp ~2015). R - f $\mathcal{B}im_N^{\text{atq}}$ category of (bicomplexes) of \mathfrak{q} -graded, free R_N -bimodules. Adjoint pair (Ad, Tr) :

$$\text{Ad}: R\text{-}f\mathcal{B}im_{N-1}^{\text{atq}} \rightarrow R\text{-}f\mathcal{B}im_N^{\text{atq}}$$

$$B \mapsto B \otimes_{R_{N-1}} (R_N^e / (x_N \otimes 1 - 1 \otimes x_N)) \quad \Leftrightarrow$$

extending scalars

$$\text{Ad} \left(\begin{array}{|c|} \hline \text{c} \\ \hline \end{array} \right) = \begin{array}{|c|} \hline \text{c} \\ \hline \end{array} \Big|$$

$$\text{Tr}: R\text{-}f\mathcal{B}im_N^{\text{atq}} \rightarrow R\text{-}f\mathcal{B}im_{N-1}^{\text{atq}}$$

$$B \mapsto (B \xrightarrow{x_N \cdot b - b \cdot x_N} \mathfrak{a}q^2 B) \quad \Leftrightarrow$$

identifying left-right action

$$\text{Tr} \left(\begin{array}{|c|} \hline \text{c} \\ \hline \end{array} \right) = \begin{array}{|c|} \hline \text{c} \\ \hline \end{array} \bigcirc$$


Skein relations. One gets e.g.

$$\begin{array}{|c|} \hline \text{D} \\ \hline \text{C} \\ \hline \text{B} \\ \hline \end{array} \bigcirc \cong \begin{array}{|c|} \hline \text{D} \\ \hline \text{C} \\ \hline \text{B} \\ \hline \end{array} \bigcirc \quad \& \quad \begin{array}{|c|} \hline 1 \\ \hline \\ \hline 1 \end{array} \bigcirc \cong \text{atq}^4 \begin{array}{|c|} \hline 1 \\ \hline \\ \hline 1 \end{array} \quad \& \quad \begin{array}{|c|} \hline 1 \\ \hline \\ \hline 1 \end{array} \bigcirc \cong \begin{array}{|c|} \hline 1 \\ \hline \\ \hline 1 \end{array}$$


Partial Hochschild homology (à la Hogancamp ~2015). R - f $\mathcal{B}im_N^{\text{atq}}$ category of (bicomplexes) of \mathfrak{q} -graded, free R_N -bimodules. Adjoint pair (Ad, Tr) :

Theorem (after normalization).

We get a triply-graded invariant $\text{HHH}_M^*(\theta) \in \mathbb{k}\text{-Vect}^{\text{atq}}$ for $\theta \in \mathcal{B}r(g, n)$, which respects Markov stabilization, *i.e.*



Skein relations. One gets e.g.



Partial Hochschild homology (à la Hogancamp ~2015). R - f $\mathcal{B}im_N^{\text{atq}}$ category of (bicomplexes) of \mathfrak{q} -graded, free R_N -bimodules. Adjoint pair (Ad, Tr) :

Theorem (after normalization).

We get a triply-graded invariant $\text{HHH}_M^*(\mathcal{C}) \in \mathbb{k}\text{-Vect}^{\text{atq}}$ for $\mathcal{C} \in \mathcal{B}r(g, n)$, which respects Markov stabilization, *i.e.*

$$\text{HHH}_M^* \left(\begin{array}{c} \text{Diagram 1} \end{array} \right) \cong \text{HHH}_M^* \left(\begin{array}{c} \text{Diagram 2} \end{array} \right) \cong \text{HHH}_M^* \left(\begin{array}{c} \text{Diagram 3} \end{array} \right)$$

Skein relations. One gets a σ

However, we are not quite there:
one gets a too strong Markov conjugation, *i.e.*

$$\text{HHH}_M^* \left(\begin{array}{c} \text{Diagram 4} \end{array} \right) \cong \text{HHH}_M^* \left(\begin{array}{c} \text{Diagram 5} \end{array} \right) \cong \text{HHH}_M^* \left(\begin{array}{c} \text{Diagram 6} \end{array} \right)$$

Partial Hochschild homology (à la Hogancamp ~2015). R - f $\mathcal{B}im_N^{\text{atq}}$ category of (bicomplexes) of \mathfrak{q} -graded, free R_N -bimodules. Adjoint pair (Ad, Tr) :

$$\text{Ad}: R\text{-}f\mathcal{B}im_{N-1}^{\text{atq}} \rightarrow R\text{-}f\mathcal{B}im_N^{\text{atq}}$$

$$\text{Ad} \left(\begin{array}{|c|} \hline \text{c} \\ \hline \end{array} \right) =$$


Idea: Flank them!

& should be thought as &

and things get stuck, e.g.

topologically stuck: & algebraically stuck:

Skein re




Partial Hochschild homology (à la Hogancamp ~2015). $R\text{-}f\mathcal{B}im_N^{\text{atq}}$ category of (bicomplexes) of \mathfrak{q} -graded, free R_N -bimodules. Adjoint pair (Ad, Tr) :


$$\text{Ad}: R\text{-}f\mathcal{B}im_{N-1}^{\text{atq}} \rightarrow R\text{-}f\mathcal{B}im_N^{\text{atq}}$$

$$B \mapsto B \otimes_{R_{N-1}} (R_N^e / (x_N \otimes 1 - 1 \otimes x_N)) \quad \longleftrightarrow \quad \text{Ad} \left(\begin{array}{|c|} \hline \text{c} \\ \hline \end{array} \right) =$$

Theorem (after normalization and flanking).

We get a triply-graded invariant $\text{HHH}_M^*(\ell) \in \mathbb{k}\text{-Vect}^{\text{atq}}$ for $\ell \in \mathcal{B}r(g, n)$, which respects Markov conjugation and stabilization, i.e.





David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16

The Markov moves on $\text{Alk}(L; \mathfrak{A}, \mathfrak{B})$ are conjugation and stabilization.

Conjugation.

$$\beta = \alpha \beta \alpha^{-1} \iff \text{for } \beta \in \text{Alk}(p, n), \alpha \in \{\beta_1, \dots, \beta_{n-1}\}$$

Stabilization.

$$(\epsilon^{-1} \beta \epsilon, \{1\}) \iff \epsilon \beta \epsilon^{-1} \cup (\epsilon^{-1} \{1\} \epsilon, \{1\}) \iff \text{for } \beta, \epsilon \in \text{Alk}(p, n)$$

They are weaker than the Markov moves.

David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16

$\text{con}(\epsilon/\delta)$ twice on a link

Currently known (to the best of my knowledge).

Genus	type A	type C
$g=0$	$\text{Alk}(n) \cong \text{AT}(A_{n-1})$	
$g=1$	$\text{Alk}(1, n) \cong \mathbb{Z} \times \text{AT}(A_{n-1}) \cong \text{AT}(A_{n-1}, 1)$	$\text{Alk}(1, n) \cong \text{AT}(C_{n-1})$
$g=2$	$\text{Alk}(2, n) \cong \text{AT}(C_{n-1})$	
$g \geq 2$		

And some $\mathbb{Z}/2\mathbb{Z}$ -orbifolds $(\mathbb{Z}/2\mathbb{Z}) \times \Sigma$ -paracore

Genus	type D	type B
$g=0$		
$g=1$	$\text{Alk}(1, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \mathbb{Z}/2\mathbb{Z} \times \text{AT}(D_{n-1})$	$\text{Alk}(1, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \text{AT}(B_{n-1})$
$g=2$	$\text{Alk}(2, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \mathbb{Z}/2\mathbb{Z} \times \text{AT}(D_{n-1})$	$\text{Alk}(2, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \mathbb{Z}/2\mathbb{Z} \times \text{AT}(B_{n-1})$
$g \geq 2$		

(For orbifolds "genus" is just an analogy)

David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16



David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16

The Markov moves on $\text{Alk}(L; \mathfrak{A}, \mathfrak{B})$ are conjugation and stabilization.

Theorem (Hiring-Olsberg-Lambropoulos – 2012).
Two links in \mathfrak{W} are equivalent if and only if they are equal in $\text{Alk}(p, n)$ up to conjugation and stabilization.

Conjugation.
for $\beta \in \mathfrak{C}$

Stabilization.
 $(\epsilon^{-1} \beta \epsilon, \{1\}) \iff \epsilon \beta \epsilon^{-1} \cup (\epsilon^{-1} \{1\} \epsilon, \{1\}) \iff \text{for } \beta, \epsilon \in \mathfrak{C}$

They are weaker than the Markov moves.

Philosophy 1: Result. Note that the type A embedding guarantees that any usual invariant of braids produces an invariant of braids in \mathfrak{W} .

same resolution


“tag” (think of $\mathbb{Z}/2\mathbb{Z}$ framing)

Genus $g = 0, 1$.

Works quite well (e.g. use “Norton-Vu”’s ideas on the categorified level)
We mimic this for M being “large, but finite”

David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16

The Alexander closure on $\text{Alk}(L; \mathfrak{A}, \mathfrak{B})$ is given by merging core strands at infinity.



wrong closure correct closure


This is different from the Alexander closure.

David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16



Figure: The Coxeter graph of finite type. (from the paper Coxeter groups and quantum groups)

Examples.
Type A_n — symmetric group
Type B_n — dihedral group
Type H_3 — dodecahedron/icosahedron — exceptional Coxeter group.
For h , we have a 4-gon:



Partial Hochschild homology (a la Hogeanscamp – 2015). $H_1 f \mathfrak{H}(\mathfrak{A}, \mathfrak{B})$ category of $(\mathbb{Z}/2\mathbb{Z}) \times \Sigma$ is graded, free \mathbb{Z}_2 -bimodules. Adjoint pair (Ad, Tr) :


$$\text{Ad}: H_1 f \mathfrak{H}(\mathfrak{A}, \mathfrak{B})^{\mathbb{Z}/2\mathbb{Z}} \rightarrow H_1 f \mathfrak{H}(\mathfrak{A}, \mathfrak{B})^{\mathbb{Z}/2\mathbb{Z}}$$

$$\text{Ad} \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) = \text{Ad} \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right)$$

We get a triply-graded invariant $\text{HHH}_1(L; \mathfrak{A}) \in \mathbb{Z}\langle \mathfrak{A}, \mathfrak{B} \rangle^{\mathbb{Z}/2\mathbb{Z}}$ for $\mathfrak{A} \in \text{Alk}(p, n)$, which respects Markov conjugation and stabilization, i.e.

David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16

There is still much to do...



David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16

The Markov moves on $\mathcal{A}(B(p, n))$ are conjugation and stabilization.

Conjugation.

$$\beta = \alpha R \alpha^{-1}$$

for $\beta \in \mathcal{A}(p, n)$, $\alpha \in \{R_1, \dots, R_{n-1}\}$

Stabilization.

$$-\epsilon \beta \cup \{e\} \cup \{e\}^{-1} \cup \beta \cup \{e\}$$

for $\beta \in \mathcal{A}(p, n)$, $\epsilon \in \{+, -\}$

They are weaker than the Markov moves.

David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16

$\text{con}(e/4)$ twice on a link

Currently known (to the best of my knowledge).


Genus	type A	type C
$g=0$	$\mathcal{A}(1, n) \cong \text{AT}(A_{n-1})$	
$g=1$	$\mathcal{A}(1, n) \cong \mathbb{Z} \times \text{AT}(A_{n-1}) \cong \text{AT}(C_{n-1})$	$\mathcal{A}(1, n) \cong \text{AT}(C_{n-1})$
$g=2$	$\mathcal{A}(1, n) \cong \mathbb{Z} \times \text{AT}(C_{n-1})$	
$g \geq 2$		

And some $\mathbb{Z}/2\mathbb{Z}$ -orbifolds $(\mathbb{Z}/2\mathbb{Z})$ -pairs:



Genus	type D	type B
$g=0$		
$g=1$	$\mathcal{A}(1, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \mathbb{Z}/2\mathbb{Z} \times \text{AT}(D_{n-1})$	$\mathcal{A}(1, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \text{AT}(B_{n-1})$
$g=2$	$\mathcal{A}(1, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \mathbb{Z}/2\mathbb{Z} \times \text{AT}(D_{n-1})$	$\mathcal{A}(1, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \mathbb{Z}/2\mathbb{Z} \times \text{AT}(B_{n-1})$
$g \geq 2$		

(For orbifolds "genus" is just an analogy.)

David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16




David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16



David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16

The Alexander closure on $\mathcal{A}(B(p, n))$ is given by merging core strands at infinity.



This is different from the Alexander closure.

David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16






Figure: The Coxeter graphs of finite type. (from the paper Coxeter graphs and root systems)

Examples. The given a generator-action presentation. Type A_n — symmetric group S_{n+1} . Type B_n — dihedral group D_{n+1} . Type H_3 — dodecahedron/icosahedron — exceptional Coxeter group. For h_n we have a 4-gon:




David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2019, 12/16

Thanks for your attention!

The Reidemeister braid relations:



These hold for usual strands only since core strands do not cross each other, e.g.




[← Back](#)

Brunn ~ 1897 , Alexander ~ 1923 . For any link ℓ in the 3-ball \mathcal{D}^3 there is a braid in $\mathcal{B}r(\infty)$ whose closure is isotopic to ℓ .

There are various proofs of this result, are all based on the same idea: “Eliminate one by one the arcs of the diagram that have the wrong sense.”


Here is an example which works in the context of general 3-manifolds: “Mark the local maxima and minima of the link diagram with respect to some height function and cut open wrong subarcs.”, e.g.




◀ Back

Markov \sim 1936. Two links in the 3-ball \mathcal{D}^3 are equivalent if and only if they are equal in $\mathcal{B}r(\infty)$ up to conjugation and stabilization.

Conjugation.




Stabilization.




[← Back](#)

Markov \sim 1936. Two links in the 3-ball \mathcal{D}^3 are equivalent if and only if they are equal in $\mathcal{B}r(\infty)$ up to conjugation and stabilization.

Conjugation.



Stabilization.







Figure: The first ever “published” braid diagram. (Page 283 from Gauß’ handwritten notes, volume seven, ≤1830).

Tits ~1961++. Gauß’ braid group is the type A case of more general groups.



Artin's approach: "Arithmetrization of braids".
 However, he still needs topological arguments.

And this is one main problem why general Artin–Tits groups are so complicated:
 Basically, they are "infinite groups without extra structure".

Ad. Gauß
cd. ob.
 Man findet mir in jeder mehr zu jeder wie oft + mit = verhält

Figure: The first ever "published" braid diagram. (Page 283 from Gauß' handwritten notes, volume seven, ≤1830).

Tits ~1961++. Gauß' braid group is the type A case of more general groups.




Figure: The Coxeter graphs of finite type. (Picture from https://en.wikipedia.org/wiki/Coxeter_group.)

Examples.


Type $A_3 \iff$ tetrahedron \iff symmetric group S_4 .

Type $B_3 \iff$ cube/octahedron \iff Weyl group $(\mathbb{Z}/2\mathbb{Z})^3 \times S_3$.

Type $H_3 \iff$ dodecahedron/icosahedron \iff exceptional Coxeter group.

For I_8 we have a 4-gon:

Idea (Coxeter \sim 1934++).






Figure: The Coxeter graphs of finite type. (Picture from https://en.wikipedia.org/wiki/Coxeter_group.)

Examples.

Type $A_3 \iff$ tetrahedron \iff symmetric group C_4


Type $B_3 \iff$ cube/octahedron \iff wreath group $(\mathbb{Z}/2\mathbb{Z}) \ltimes S_3$.

Type $H_3 \iff$ dodecahedron/icosahedron \iff exceptional Coxeter group.

For I_8 we have a 4-gon:

Fix a flag F .

Idea (Coxeter $\sim 1934++$).






Figure: The Coxeter graphs of finite type. (Picture from https://en.wikipedia.org/wiki/Coxeter_group.)

Examples.

Type $A_3 \iff$ tetrahedron \iff symmetric group S_4 .

Type $B_3 \iff$ cube/octahedron \iff Weyl group $(\mathbb{Z}/2\mathbb{Z})^3 \times S_3$.


Type $H_3 \iff$ dodecahedron/icosahedron \iff exceptional Coxeter group.

For I_8 we have a 4-gon:

Fix a flag F .

Idea (Coxeter $\sim 1934++$).

Fix a hyperplane H_0 permuting the adjacent 0-cells of F .






Figure: The Coxeter graphs of finite type. (Picture from https://en.wikipedia.org/wiki/Coxeter_group.)

Examples.

Type $A_3 \iff$ tetrahedron \iff symmetric group S_4 .

Type $B_3 \iff$ cube/octahedron \iff Weyl group $(\mathbb{Z}/2\mathbb{Z})^3 \times S_3$.

Type $H_3 \iff$ dodecahedron/icosahedron \iff exceptional Coxeter group.


For I_8 we have a 4-gon:

Fix a flag F .

Idea (Coxeter \sim 1934++).

Fix a hyperplane H_0 permuting the adjacent 0-cells of F .

Fix a hyperplane H_1 permuting the adjacent 1-cells of F , etc.






Figure: The Coxeter graphs of finite type. (Picture from https://en.wikipedia.org/wiki/Coxeter_group.)

Examples.

Type $A_3 \iff$ tetrahedron \iff symmetric group S_4 .

Type $B_3 \iff$ cube/octahedron \iff Weyl group $(\mathbb{Z}/2\mathbb{Z})^3 \times S_3$.

Type $H_3 \iff$ dodecahedron/icosahedron \iff exceptional Coxeter group.

For I_8 we have a 4-gon:


Fix a flag F .

Idea (Coxeter $\sim 1934++$).

Fix a hyperplane H_0 permuting the adjacent 0-cells of F .

Fix a hyperplane H_1 permuting the adjacent 1-cells of F , etc.

Write a vertex i for each H_i .






Figure: The Coxeter graphs of finite type. (Picture from https://en.wikipedia.org/wiki/Coxeter_group.)

Examples.

This gives a generator-relation presentation.

Type $A_3 \iff$ tetrahedron \iff symmetric group S_4 .

Type $B_3 \iff$ And the braid relation measures the angle between hyperplanes.

Type $H_3 \iff$ dodecahedron/icosahedron \iff exceptional Coxeter group.

For I_8 we have a 4-gon:

Fix a flag F .


Idea (Coxeter $\sim 1934++$).

Fix a hyperplane H_0 permuting the adjacent 0-cells of F .

Fix a hyperplane H_1 permuting the adjacent 1-cells of F , etc.

Write a vertex i for each H_i .

Connect i, j by an n -edge for H_i, H_j having angle $\cos(\pi/n)$.



Three gradings:

$\mathfrak{q} \leftrightarrow$ internal

&

$\mathfrak{t} \leftrightarrow$ homological

&

$\mathfrak{a} \leftrightarrow$ Hochschild

Example. To compute Hochschild cohomology take the Koszul resolution

$$\bigotimes_{i=1}^N \left(R^e = R \otimes R^{\text{op}} \xrightarrow{\cdot(x_i \otimes 1 - 1 \otimes x_i)} \mathfrak{a} \mathfrak{q}^2 R^e \right),$$

Tensor it with B , gives a complex with differentials $x_i \otimes 1 - 1 \otimes x_i$, of which we think as identifying the variables. This gives a chain complex having non-trivial chain groups in \mathfrak{a} -degree $0, \dots, n$. Here the i^{th} chain group consists of $\binom{n}{i}$ copies of B , with differentials given by the various ways of identifying i variables. The a^{th} cohomology = a^{th} Hochschild cohomology.

Example. If B is already a \mathfrak{t} -graded complex, then one can take homology of it and gets “triple H ”.