


# HOMFLYPT homology for links in handlebodies


Or: All I know about Artin–Tits groups; and a filler for the remaining 49 minutes


Daniel Tubbenhauer





Joint with David Rose


February 2019










































































## 1 Links and braids in handlebodies

- Braid diagrams
- Links in handlebodies

## 2 Some “low-genus-coincidences”


- The ball and the torus
- The torus and the double torus

## 3 Arbitrary genus


- Braid invariants – some ideas
- Link invariants – some ideas

Let  $\text{Br}(g, n)$  be the group defined as follows.


**Generators.** Braid and twist generators



**Relations.** [Reidemeister braid relations](#), type C relations and special relations, e.g.





Involves three players and inverses!



Let  $\text{Br}(g, n)$  be the group defined as follows.

**Generators.** Braid and twist

$\sigma_i \leftrightarrow$




**Relations.**

► Reidemeister braid relation


and special relations, e.g.

Involves three players and inverses!



$$\sigma_1 t_2 \sigma_1 t_2 = t_2 \sigma_1 t_2 \sigma_1$$

&




$$(\sigma_1 t_2 \sigma_1^{-1}) t_3 = t_3 (\sigma_1 t_2 \sigma_1^{-1})$$


Let  $\text{Br}(g, n)$  be the group defined as follows.

**Generators.** Braid and twist

$$\sigma_i \leftrightarrow$$



**Example.**



**Relations.**



► Reidemeister braid relation

and special relations, e.g.

Involves three players and inverses!



=



$$\sigma_1 t_2 \sigma_1 t_2 = t_2 \sigma_1 t_2 \sigma_1$$

$$(\sigma_1 t_2 \sigma_1^{-1}) t_3 = t_3 (\sigma_1 t_2 \sigma_1^{-1})$$

Let  $\text{Br}(g, n)$  be the group defined as follows.

**Generators.** Braid and twist generators

1 g 1 i i+1 n                      1 i g 1 2 n

**Fact (type A embedding).**

$\text{Br}(g, n)$  is a subgroup of the usual braid group  $\mathcal{B}\text{r}(g+n)$ .

Relatio


A visualization exercise.

$\ell_1 t_2 \ell_1^{-1} t_2 = t_2 \ell_1 t_2 \ell_1^{-1}$                        $(\ell_1 t_2 \ell_1^{-1}) t_3 = t_3 (\ell_1 t_2 \ell_1^{-1})$

The group  $\mathcal{B}r(g, n)$  of braid in a  $g$ -times punctures disk  $\mathcal{D}_g^2 \times [0, 1]$ :


---

Two types of braidings, the usual ones and “winding around cores”, e.g.



Theorem (Häring-Oldenburg–Lambropoulou ~2002, Vershinin ~1998).


The map



is an isomorphism of groups  $Br(g, n) \rightarrow \mathcal{B}r(g, n)$ .

The group  $\mathcal{B}r(g, n)$  of braid in a  $g$ -times punctures disk  $\mathcal{D}_g^2 \times [0, 1]$ :

Two types of embeddings, e.g.




The group  $\mathcal{B}r(g, n)$  of braid in a  $g$ -times punctures disk  $\mathcal{D}_g^2 \times [0, 1]$ :

Two types of braidings, the usual ones and “winding around cores” e.g.

**Note.**

For the proof it is crucial that  $\mathcal{D}_g^2$  and the boundary points of the braids  $\bullet$  are only defined up to isotopy, e.g.




$\Rightarrow$  one can always “conjugate cores to the left”.


This is useful to define  $\mathcal{B}r(g, \infty)$ .

The Alexander closure on  $\mathcal{B}r(g, \infty)$  is given by merging core strands at infinity.

---



wrong closure




correct closure

This is different from the [classical](#) Alexander closure.

The Alexander closure on  $\mathcal{B}r(g, \infty)$  is given by merging core strands at infinity.

**Theorem (Lambropoulou ~1993).**

For any link  $\ell$  in the genus  $g$  handlebody  $\mathcal{H}_g$  there is a braid in  $\mathcal{B}r(g, \infty)$  whose (correct!) closure is isotopic to  $\ell$ .



This is different from the [classical](#) Alexander closure.


The Alexander closure on  $\mathcal{B}r(g, \infty)$  is given by merging core strands at infinity.

**Theorem (Lambropoulou ~1993).**


For any link  $\ell$  in the genus  $g$  handlebody  $\mathcal{H}_g$  there is a braid in  $\mathcal{B}r(g, \infty)$  whose (correct!) closure is isotopic to  $\ell$ .

**Fact.**


$\mathcal{H}_g$  is given by a complement in the 3-sphere  $\mathcal{S}^3$  by an open tubular neighborhood of the embedded graph obtained by gluing  $g + 1$  unknotted "core" edges to two vertices.



the 3-ball  $\mathcal{H}_0 = \mathcal{D}^3$



a torus  $\mathcal{H}_1$



$\mathcal{H}_2$


This is

The Markov moves on  $\mathcal{B}r(g, \infty)$  are conjugation and stabilization.

## Conjugation.

$$\ell \sim s\ell s^{-1}$$

for  $\ell \in \mathcal{B}r(g, n)$ ,  $s \in \langle \ell_1, \dots, \ell_{n-1} \rangle$




## Stabilization.

$$(c\uparrow)\ell_n(\ell\uparrow)$$

$$\sim c\ell \sim (c\uparrow)\ell_n^{-1}(\ell\uparrow)$$

for  $\ell, c \in \mathcal{B}r(g, n)$ ,




They are weaker than the classical Markov moves.

**Theorem (Häring-Oldenburg–Lambropoulou ~2002).**

Two links in  $\mathcal{H}_g$  are equivalent if and only if they are equal in  $\mathcal{B}r(g, \infty)$  up to conjugation and stabilization.

$$\ell \sim s\ell s^{-1}$$

for  $\ell \in \mathcal{B}r(g, n)$ ,  $s \in \langle \ell_1, \dots, \ell_{n-1} \rangle$




**Stabilization.**

$$(c\uparrow)\ell_n(\ell\uparrow)$$

$$\sim c\ell \sim (c\uparrow)\ell_n^{-1}(\ell\uparrow)$$

for  $\ell, c \in \mathcal{B}r(g, n)$ ,





They are weaker than the classical Markov moves.

The Markov moves on  $\mathcal{B}r(g, \infty)$  are conjugation and stabilization

**Theorem (Häring-Oldenburg–Lambropoulou ~2002).**

Two links in  $\mathcal{H}_g$  are equivalent if and only if they are equal in  $\mathcal{B}r(g, \infty)$  up to conjugation and stabilization.

**Example.**



for  $\beta \in \mathcal{B}r$

Stabilization

$(c\uparrow)\beta$

$\sim c\beta \sim (c\downarrow)\beta$


for  $\beta, c \in \mathcal{B}r$

They are weakly

The Markov moves on  $\mathcal{B}r(g, \infty)$  are conjugation and stabilization.


## Conjugation.

$\ell \sim s\ell s^{-1}$   
for  $\ell \in \mathcal{B}r(g, n)$ ,



## Stabilization.

$(c\uparrow)\ell_n(\ell\uparrow)$   
 $\sim c\ell \sim (c\uparrow)\ell_n^{-1}(\ell\uparrow)$   
for  $\ell, c \in \mathcal{B}r(g, n)$ ,




They are weaker than the classical Markov moves.

The Markov moves on  $\mathcal{B}r(g, \infty)$  are conjugation and stabilization.

## Conjugation.


$\ell \sim s\ell s^{-1}$   
for  $\ell \in \mathcal{B}r(g, n)$ ,



## Stabilization.

$(c\uparrow)\ell_n(\ell\uparrow)$   
 $\sim c\ell \sim (c\uparrow)\ell_n^{-1}(\ell\uparrow)$   
for  $\ell, c \in \mathcal{B}r(g, n)$ ,

Let me explain what we can do.



They are weaker than the classical Markov moves.

Let  $\Gamma$  be a Coxeter graph.

---

**Artin**  $\sim$ 1925, **Tits**  $\sim$ 1961++. The Artin–Tits group and its Coxeter group quotient are given by generators-relations:

$$\begin{aligned} \text{AT}(\Gamma) &= \langle \ell_i \mid \underbrace{\cdots \ell_i \ell_j \ell_i}_{m_{ij} \text{ factors}} = \underbrace{\cdots \ell_j \ell_i \ell_j}_{m_{ij} \text{ factors}} \rangle \\ &\downarrow \\ \text{W}(\Gamma) &= \langle \sigma_i \mid \sigma_i^2 = 1, \underbrace{\cdots \sigma_i \sigma_j \sigma_i}_{m_{ij} \text{ factors}} = \underbrace{\cdots \sigma_j \sigma_i \sigma_j}_{m_{ij} \text{ factors}} \rangle \end{aligned}$$

Artin–Tits groups [▶ generalize](#) classical braid groups, Coxeter groups [▶ generalize](#) polyhedron groups.

$\cos(\pi/3)$  on a line:

type  $A_{n-1}$ :  $1 \text{ --- } 2 \text{ --- } \dots \text{ --- } n-2 \text{ --- } n-1$

---

**The classical case.** Consider the map

$$\beta_i \mapsto \begin{array}{cccc} 1 & i & i+1 & n \\ \uparrow & \nearrow & \nearrow & \uparrow \\ \dots & \times & \dots & \\ \uparrow & \searrow & \searrow & \uparrow \\ 1 & i & i+1 & n \end{array} \quad \text{braid rel.:} \quad \begin{array}{c} \uparrow \uparrow \uparrow \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \uparrow \uparrow \uparrow \end{array} = \begin{array}{c} \uparrow \uparrow \uparrow \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \uparrow \uparrow \uparrow \end{array}$$

---

**Artin ~1925.** This gives an isomorphism of groups  $AT(A_{n-1}) \xrightarrow{\cong} \mathcal{B}r(0, n)$ .

$\cos(\pi/3)$  on a line:

Jones ~1987.

Markov trace on the Hecke algebra of type A

$\rightsquigarrow$  two variable  $q, a$  polynomial invariant (HOMFLYPT polynomial).

The clas

$q$ =Hecke parameter ;  $a$ =trace parameter .



braid rel.:



Artin ~1925. This gives an isomorphism of groups  $AT(A_{n-1}) \xrightarrow{\cong} \mathcal{B}r(0, n)$ .

I will come back to this with more details for general genus  $g$ .  
For the time being: This works quite well!

$\cos(\pi/3)$  on a line:

**Jones ~1987.**

Markov trace on the Hecke algebra of type A

$\rightsquigarrow$  two variable  $q, a$  polynomial invariant (HOMFLYPT polynomial).

The clas

$q$ =Hecke parameter ;  $a$ =trace parameter .

**Khovanov ~2005; categorification.**

Hochschild homology on complexes of the Hecke category of type A

$\rightsquigarrow$  "three variable  $q, t, a$  homological invariant" (HOMFLYPT homology).

$q$ =Hecke parameter ;  $t$ =homological parameter ;  $a$ =Hochschild parameter .


Artin ~1929. This gives an isomorphism of groups  $AI(A_{n-1}) \cong \mathcal{B}I(0, n)$ .

I will come back to this with more details for general genus  $g$ .  
For the time being: This works quite well!

$\cos(\pi/3)$  on a circle.



**Affine adds genus.** Consider the map



**tom Dieck ~1998. (Earlier reference?)** This gives an isomorphism of groups  $\mathbb{Z} \times \text{AT}(\tilde{A}_{n-1}) \xrightarrow{\cong} \mathcal{B}r(1, n)$ .

$\cos(\pi/3)$  on a circle.


**Orellana–Ram ~2004. (Earlier reference?)**

Markov trace on the Hecke algebra of type  $\tilde{A}$

Affine  $a$

$\rightsquigarrow$  two variable  $q, a$  polynomial invariant (HOMFLYPT polynomial).

$q$ =Hecke parameter ;  $a$ =trace parameter .



$\beta_0 \mapsto$

$\&$   $\beta_i \mapsto$



braid rel.:



**tom Dieck ~1998. (Earlier reference?)** This gives an isomorphism of groups

$\mathbb{Z} \times \text{AT}(\tilde{A}_{n-1}) \cong \mathcal{Rr}(1, n)$

I will come back to this with more details for general genus  $g$ .  
For the time being: This works quite well!

$\cos(\pi/3)$  on a circle.

**Orellana–Ram ~2004. (Earlier reference?)**

Markov trace on the Hecke algebra of type  $\tilde{A}$

↪ two variable  $q, a$  polynomial invariant (HOMFLYPT polynomial).

$q$ =Hecke parameter ;  $a$ =trace parameter .

Affine a

**???; categorification.**

Hochschild homology on complexes of the Hecke category of type  $\tilde{A}$

↪ “three variable  $q, t, a$  homological invariant” (HOMFLYPT homology).

$q$ =Hecke parameter ;  $t$ =homological parameter ;  $a$ =Hochschild parameter .

tom Dieck ~1996. (Earlier reference?) This gives an isomorphism of groups

$\mathbb{Z} \times \text{AT}(\tilde{A}_{n-1}) \cong \mathcal{R}(1, n)$

I will come back to this with more details for general genus  $g$ .  
For the time being: This works quite well!


$\cos(\pi/3)$  on a circle.

**Fact.** One can recover the (missing) generator of  $\mathbb{Z}$  if one works with extended affine type A.


Affine adds

$\beta_0 \mapsto$


"extended, extra generator"  $\mapsto$



and



give



tom Dieck


$$\mathbb{Z} \times \text{AT}(\tilde{A}_{n-1}) \xrightarrow{\cong} \mathcal{B}r(1, n).$$

m of groups

$\cos(\pi/4)$  on a line:

$$\text{type } C_n: 0 \stackrel{4}{=} 1 - 2 - \dots - n-1 - n$$

**The semi-classical case.** Consider the map



**Brieskorn  $\sim$  1973.** This gives an isomorphism of groups  $\text{AT}(C_n) \xrightarrow{\cong} \mathcal{B}r(1, n)$ .


$\cos(\pi/4)$  on a line:

**Geck–Lambropoulou ~1997.**

Markov trace on the Hecke algebra of type C

$\rightsquigarrow$  two variable  $q, a$  polynomial invariant (HOMFLYPT polynomial).

$q$ =Hecke parameter ;  $a$ =trace parameter .



**Brieskorn ~1973.** This gives an isomorphism of groups  $AT(C_n) \xrightarrow{\cong} \mathcal{B}r(1, n)$ .

I will come back to this with more details for general genus  $g$ .  
For the time being: This works quite well!

$\cos(\pi/4)$  on a line:

**Geck–Lambropoulou ~1997.**

Markov trace on the Hecke algebra of type C

$\rightsquigarrow$  two variable  $q$ , a polynomial invariant (HOMFLYPT polynomial).

$q$ =Hecke parameter ;  $a$ =trace parameter .

1 2 n 1 i i+1 n

**Rouquier ~2012, Webster–Williamson ~2009; categorification.**

Hochschild homology on complexes of the Hecke category of type C

$\rightsquigarrow$  “three variable  $q$ ,  $t$ ,  $a$  homological invariant” (HOMFLYPT homology).

$q$ =Hecke parameter ;  $t$ =homological parameter ;  $a$ =Hochschild parameter .

Brieskorn ~1975. This gives an isomorphism of groups  $A_1(\mathbb{C}_n) \rightarrow \mathcal{S}_1(1, n)$ .

I will come back to this with more details for general genus  $g$ .  
For the time being: This works quite well!



$\beta_0$

Brieskorn

$\cos(\pi/4)$  on a line:

**Fact. (Not true in type A.)**

There is a whole infinite family of Markov traces,  
one for each choice of a value for essential unlinks.

The

$\beta_0$

extra parameter and extra parameter etc.

However, I only know the categorification of one of these.

**Brieskorn  $\sim$ 1973.** This gives an isomorphism of groups  $AT(C_n) \xrightarrow{\cong} \mathcal{B}r(1, n)$ .

$\cos(\pi/4)$  on a line:

**Fact. (Not true in type A.)**

There is a whole infinite family of Markov traces,  
one for each choice of a value for essential unlinks.

The

$\beta_0$

extra parameter and extra parameter etc.

However, I only know the categorification of one of these.

**Fact. (Not true in type A.)**

Brieskorn  $\sim$ 1973. There is also a second Hecke parameter,  $\mathbb{C}[n] \xrightarrow{\mathbb{R}} \mathcal{B}r(1, n)$ ,  
which we do not know how to categorify yet.

$\cos(\pi/4)$  twice on a line:

$$\text{type } \tilde{C}_n: 0^1 \stackrel{4}{=} 1 \text{ --- } 2 \text{ --- } \dots \text{ --- } n-1 \text{ --- } n \stackrel{4}{=} 0^2$$


**Affine adds genus.** Consider the map

$$\beta_{0^1} \mapsto \begin{array}{c} \color{red}{1} \quad 1 \quad n \quad \color{red}{2} \\ \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \\ \color{red}{1} \quad 1 \quad n \quad \color{red}{2} \end{array} \quad \& \quad \beta_i \mapsto \begin{array}{c} i \quad i+1 \\ \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \\ i \quad i+1 \end{array} \quad \& \quad \beta_{0^2} \mapsto \begin{array}{c} \color{red}{1} \quad 1 \quad n \quad \color{red}{2} \\ \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \quad \color{red}{\rule{0.2em}{1em}} \\ \color{red}{1} \quad 1 \quad n \quad \color{red}{2} \end{array}$$


**Allcock ~1999.** This gives an isomorphism of groups  $\text{AT}(\tilde{C}_n) \xrightarrow{\cong} \mathcal{B}r(2, n)$ .

$\cos(\pi/4)$  twice


This case is strange – it only arises under conjugation:



By a miracle, one can avoid the special relation



This relation involves three players and inverses. Bad!




Affine adds ge

$\beta_{01} \mapsto$

Allcock  $\sim$ 1999. This gives an isomorphism of groups  $AT(\tilde{C}_n) \xrightarrow{\cong} \mathcal{B}r(2, n)$ .

This case is strange – it only arises under conjugation:


$\cos(\pi/4)$  twice



Affine adds ge


By a miracle, one can avoid the special relation

$\beta_{0^1} \mapsto$



This relation involves three players and inverses.

Bad!




Currently, not much seems to be known, but I think the same story works.


Allcock  $\sim$ 1999. This gives an isomorphism of groups  $AT(\tilde{C}_n) \xrightarrow{\cong} \mathcal{B}r(2, n)$ .

$\cos(\pi/4)$  twice

This case is strange – it only arises under conjugation:



By a miracle, one can avoid the special relation




This relation involves three players and inverses. Bad!



Currently, not much seems to be known, but I think the same story works.

Allcock However, this is where it seems to end, e.g. genus  $g = 3$  wants to be  $n$ ).



But the special relation makes it a mere quotient.  
So: In the remaining time I tell you what works.

$\cos(\pi/4)$  twice on a line:

Currently known (to the best of my knowledge).

Aff

Genus	type A	type C
$g = 0$	$\mathcal{B}r(n) \cong AT(A_{n-1})$	
$g = 1$	$\mathcal{B}r(1, n) \cong \mathbb{Z} \ltimes AT(\tilde{A}_{n-1}) \cong AT(\hat{A}_{n-1})$	$\mathcal{B}r(1, n) \cong AT(C_n)$
$g = 2$		$\mathcal{B}r(2, n) \cong AT(\tilde{C}_n)$
$g \geq 3$		

And some  $\mathbb{Z}/2\mathbb{Z}$ -orbifolds ( $\mathbb{Z}/\infty\mathbb{Z}$  = puncture):

All


Genus	type D	type B
$g = 0$		
$g = 1$	$\mathcal{B}r(1, n)_{\mathbb{Z}/2\mathbb{Z}} \cong AT(D_n)$	$\mathcal{B}r(1, n)_{\mathbb{Z}/\infty\mathbb{Z}} \cong AT(B_n)$
$g = 2$	$\mathcal{B}r(2, n)_{\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}} \cong AT(\tilde{D}_n)$	$\mathcal{B}r(2, n)_{\mathbb{Z}/\infty\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}} \cong AT(\tilde{B}_n)$
$g \geq 3$		

(For orbifolds "genus" is just an analogy.)




# Philosophy 1: Reshetikhin–Turaev with “huge” colors.

---




# Philosophy 1: Reshetikhin–Turaev with “huge” colors.


---




# Philosophy 1: Reshetikhin–Turaev with “huge” colors.




# Philosophy 1: Reshetikhin–Turaev with “huge” colors.




# Philosophy 1: Reshetikhin–Turaev with “huge” colors.




# Philosophy 1: Reshetikhin–Turaev with “huge” colors.




# Philosophy 1: Reshetikhin–Turaev with “huge” colors.



# Philosophy 1: Reshetikhin–Turaev with “huge” colors.




# Philosophy 1: Reshetikhin–Turaev with “huge” colors.




# Philosophy 1: Resh

Note that the type A embedding guarantees that any usual invariant of braids produces an invariant of braids in  $\mathcal{H}_g$ .




Note that the type A embedding guarantees that any usual invariant of braids produces an invariant of braids in  $\mathcal{H}_g$ .



Genus  $g = 0, 1$ .

Works quite well (e.g. use Naisse–Vaz's ideas on the categorified level).

Note that the type A embedding guarantees that any usual invariant of braids produces an invariant of braids in  $\mathcal{H}_g$ .



Genus  $g = 0, 1$ .

Works quite well (e.g. use Naisse–Vaz’s ideas on the categorified level).

We mimic this for M being “huge, but finite”.

## Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$ .

---

Tuples  $\mathbf{I} = (k_1, \dots, k_N) \in \mathbb{N}_{\geq 1}^N$  with  $k_1 + \dots + k_N = N \iff$  parabolic subgroups

$$W_{\mathbf{I}} = W(A_{k_1-1}) \times \dots \times W(A_{k_N-1}) \subset W.$$

$W$  acts on  $\mathbb{R} = \mathbb{R}_N = \mathbb{k}[x_1, \dots, x_N]$  via permutation  $\rightsquigarrow$  rings of invariants  $\mathbb{R}^{\mathbf{I}}$ .

---

**Bimodules.** Identities, restriction (“merge”) and induction (“split”), e.g.

$$\begin{array}{c} 1 & 1 & 1 \\ | & | & | \\ 1 & 1 & 1 \end{array} \iff \mathbb{R}^{(1,1,1)} = \mathbb{R}, \quad \begin{array}{c} 2 & 1 \\ | & | \\ 2 & 1 \end{array} \iff \mathbb{R}^{(2,1)} = \mathbb{R}^{\sigma_1} = \mathbb{k}[x_1 + x_2, x_1 x_2, x_3].$$

$$\begin{array}{c} k+l \\ \cup \\ k \quad l \end{array} \iff \text{shift} \mathbb{R}^{(k+l)} \otimes_{\mathbb{R}^{(k+l)}} \mathbb{R}^{(k,l)}, \quad \begin{array}{c} k \quad l \\ \cup \\ k+l \end{array} \iff \mathbb{R}^{(k,l)} \otimes_{\mathbb{R}^{(k+l)}} \mathbb{R}^{(k+l)}.$$

---

Define  $\mathcal{S}_s^q(W)$  as the full 2-subcategory of the rings&bimodules 2-category.

# Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$ .

---

Tuples  $\mathbf{I} = (k_1, \dots, k_N) \in \mathbb{N}_{\geq 1}^N$  with  $k_1 + \dots + k_N = N \iff$  parabolic subgroups

$$W_{\mathbf{I}} = W(A_{k_1-1}) \times \dots \times W(A_{k_N-1}) \subset W.$$

$W$  acts on  $\mathbb{R} = \mathbb{R}_N = \mathbb{R}[x_1, \dots, x_N]$ . Rings of invariants  $\mathbb{R}^{\mathbf{I}}$ .

Everything is  $\mathbb{Z}$ -graded, called  $\mathfrak{q}$ -grading.  
I just omit this for simplicity.

**Bimodules.** Identities, restriction (“merge”) and induction (“split”), e.g.

$$\begin{array}{c} 1 & 1 & 1 \\ | & | & | \\ 1 & 1 & 1 \end{array} \iff \mathbb{R}^{(1,1,1)} = \mathbb{R}, \quad \begin{array}{c} 2 & 1 \\ | & | \\ 2 & 1 \end{array} \iff \mathbb{R}^{(2,1)} = \mathbb{R}^{\sigma_1} = \mathbb{k}[x_1 + x_2, x_1 x_2, x_3].$$

$$\begin{array}{c} k+l \\ \cup \\ k \quad l \end{array} \iff \text{shift} \mathbb{R}^{(k+l)} \otimes_{\mathbb{R}^{(k+l)}} \mathbb{R}^{(k,l)}, \quad \begin{array}{c} k \quad l \\ \cup \\ k+l \end{array} \iff \mathbb{R}^{(k,l)} \otimes_{\mathbb{R}^{(k+l)}} \mathbb{R}^{(k+l)}.$$

Define  $\mathcal{S}_s^q(W)$  as the full 2-subcategory of the rings&bimodules 2-category.

# Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$ .

A monoidal structure is given by

$$\begin{array}{c} 1 & 1 \\ \cup & \\ 1 & 1 \end{array} = \begin{array}{c} 2 \\ | \\ 1 & 1 \end{array} \leftarrow \text{glue} \rightarrow \begin{array}{c} 1 & 1 \\ \cup & \\ & 2 \end{array} \iff R \otimes_{R^{\sigma_1}} R \cong R \otimes_{R^{\sigma_1}} R^{\sigma_1} \otimes_{R^{\sigma_1}} R.$$

This gives a way to define bimodules associated to any web built out of merge and split.

**Bimodules.** Identities, restriction (“merge”) and induction (“split”), e.g.

$$\begin{array}{c} 1 & 1 & 1 \\ | & | & | \\ 1 & 1 & 1 \end{array} \iff R^{(1,1,1)} = R, \quad \begin{array}{c} 2 & 1 \\ | & | \\ 2 & 1 \end{array} \iff R^{(2,1)} = R^{\sigma_1} = \mathbb{k}[x_1 + x_2, x_1 x_2, x_3].$$

$$\begin{array}{c} k+l \\ \cup \\ k & l \end{array} \iff \text{shift} R^{(k+l)} \otimes_{R^{(k+l)}} R^{(k,l)}, \quad \begin{array}{c} k & l \\ \cup \\ k+l \end{array} \iff R^{(k,l)} \otimes_{R^{(k+l)}} R^{(k+l)}.$$

Define  $\mathcal{S}_s^q(W)$  as the full 2-subcategory of the rings&bimodules 2-category.



# Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$ .

Soergel ~1992, Williamson ~2010.

Tuples  $I = (i_1, \dots, i_N) \in \{1, \dots, N\}^N$  categorifies the Hecke algebra (or rather, the algebroid). subgroups

$$W_I = W(A_{k_1-1}) \times \cdots \times W(A_{k_N-1}) \subset W.$$

$W$  acts on  $R = R_N = \mathbb{k}[x_1, \dots, x_N]$  via permutation  $\rightsquigarrow$  rings of invariants  $R^I$ .

**Bimodules.** Identities, restriction (“merge”) and induction (“split”), e.g.

$$\begin{array}{c} 1 & 1 & 1 \\ | & | & | \\ 1 & 1 & 1 \end{array} \rightsquigarrow R^{(1,1,1)} = R, \quad \begin{array}{c} 2 & 1 \\ | & | \\ 2 & 1 \end{array} \rightsquigarrow R^{(2,1)} = R^{\sigma_1} = \mathbb{k}[x_1 + x_2, x_1 x_2, x_3].$$

$$\begin{array}{c} k+l \\ \cup \\ k \quad l \end{array} \rightsquigarrow \text{shift} R^{(k+l)} \otimes_{R^{(k+l)}} R^{(k,l)}, \quad \begin{array}{c} k \quad l \\ \cup \\ k+l \end{array} \rightsquigarrow R^{(k,l)} \otimes_{R^{(k+l)}} R^{(k+l)}.$$

Define  $\mathcal{S}_s^q(W)$  as the full 2-subcategory of the rings&bimodules 2-category.

# Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$ .

Soergel ~1992, Williamson ~2010.

Tuples  $I = (i_1, \dots, i_n) \in \{1, \dots, N\}^n$  categorifies the Hecke algebra (or rather, the algebroid) of subgroups

Rouquier ~2004, Mackaay–Stošić–Vaz ~2008, Webster–Williamson ~2009, etc.

There are certain complex (“t-graded”) of singular Soergel bimodules, e.g.

$$[\beta_i]_M = \begin{array}{c} l \quad k \\ \diagdown \quad \diagup \\ k \quad l \end{array} = \begin{array}{c} k-l \\ \diagdown \quad \diagup \\ k \quad l \\ 0 \end{array} \xrightarrow{d_0^+} \mathbf{qt} \begin{array}{c} k-l \\ \diagdown \quad \diagup \\ k \quad l \\ 1 \end{array} \xrightarrow{d_1^+} \dots \xrightarrow{d_{l-1}^+} \mathbf{q}^l \mathbf{t}^l \begin{array}{c} k \\ \diagdown \quad \diagup \\ k \quad l \end{array}$$

providing a categorical action of the Artin–Tits group of type A.

1 1 1

2 1



$$\iff \text{shift} R^{(k+l)} \otimes_{R^{(k+l)}} R^{(k,l)},$$



$$\iff R^{(k,l)} \otimes_{R^{(k+l)}} R^{(k+l)}.$$

Define  $\mathcal{S}_s^q(W)$  as the full 2-subcategory of the rings&bimodules 2-category.

# Singular Soergel bimodules $\mathcal{S}_s^q(W)$ for $W = W(A_{N-1})$ .

Soergel ~1992, Williamson ~2010.

Tuples  $\Gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{Z}^n$  categorifies the Hecke algebra (or rather, the algebroid) of subgroups

Rouquier ~2004, Mackaay–Stošić–Vaz ~2008, Webster–Williamson ~2009, etc.


There are certain complex (“t-graded”) of singular Soergel bimodules, e.g.

$$[[\beta_i]]_M = \begin{array}{c} l \quad k \\ \diagdown \quad \diagup \\ k \quad l \end{array} = \begin{array}{c} k-l \\ \diagdown \quad \diagup \\ k \quad l \\ 0 \end{array} \xrightarrow{d_0^+} \mathbf{qt} \begin{array}{c} k-l \\ \diagdown \quad \diagup \\ k \quad l \\ +1 \\ 1 \end{array} \xrightarrow{d_1^+} \dots \xrightarrow{d_{l-1}^+} \mathbf{q}^l \mathbf{t}^l \begin{array}{c} k \\ \diagdown \quad \diagup \\ k \quad l \\ l \end{array}$$

providing a categorical action of the Artin–Tits group of type A.

1 1 1 2 1

Hence, we are in business by taking  $M \gg n$ :



$R^{(k+l)}$ .

Define  $\mathcal{S}_s^q(\mathfrak{b})$  **Fact.** This gives a faithful invariant of  $[[\mathfrak{b}]]_M$  of  $\mathfrak{b} \in \mathcal{B}\mathfrak{r}(g, n)$ .



Partial Hochschild homology (à la Hogancamp ~2015).  $R$ - $f$   $\mathcal{B}im_N^{\text{atq}}$  category of (bicomplexes) of  $\mathfrak{q}$ -graded, free  $R_N$ -bimodules. Adjoint pair  $(\text{Ad}, \text{Tr})$ :

**Theorem (after normalization).**

We get a triply-graded invariant  $\text{HHH}_M^*(\mathcal{C}) \in \mathbb{k}\text{-Vect}^{\text{atq}}$  for  $\mathcal{C} \in \mathcal{B}r(g, n)$ , which respects Markov stabilization, *i.e.*

$$\text{HHH}_M^* \left( \begin{array}{c} \text{c} \\ \text{b} \end{array} \right) \cong \text{HHH}_M^* \left( \begin{array}{c} \text{c} \\ \text{b} \end{array} \right) \cong \text{HHH}_M^* \left( \begin{array}{c} \text{c} \\ \text{b} \end{array} \right)$$

**Skein relations.** One gets *e.g.*

Partial Hochschild homology (à la Hogancamp ~2015).  $R$ - $f$   $\mathcal{B}im_N^{\text{atq}}$  category of (bicomplexes) of  $\mathfrak{q}$ -graded, free  $R_N$ -bimodules. Adjoint pair  $(\text{Ad}, \text{Tr})$ :

### Theorem (after normalization).

We get a triply-graded invariant  $\text{HHH}_M^*(\mathcal{C}) \in \mathbb{k}\text{-Vect}^{\text{atq}}$  for  $\mathcal{C} \in \mathcal{B}r(g, n)$ , which respects Markov stabilization, *i.e.*

$$\text{HHH}_M^* \left( \begin{array}{c} \text{Diagram 1} \end{array} \right) \cong \text{HHH}_M^* \left( \begin{array}{c} \text{Diagram 2} \end{array} \right) \cong \text{HHH}_M^* \left( \begin{array}{c} \text{Diagram 3} \end{array} \right)$$

Skein relations. One gets a  $\sigma$

However, we are not quite there:  
one gets a too strong Markov conjugation, *i.e.*

$$\text{HHH}_M^* \left( \begin{array}{c} \text{Diagram 4} \end{array} \right) \cong \text{HHH}_M^* \left( \begin{array}{c} \text{Diagram 5} \end{array} \right) \cong \text{HHH}_M^* \left( \begin{array}{c} \text{Diagram 6} \end{array} \right)$$

Partial Hochschild homology (à la Hogancamp ~2015).  $R\text{-}f\mathcal{B}im_N^{\text{at}q}$  category of (bicomplexes) of  $q$ -graded, free  $R_N$ -bimodules. Adjoint pair  $(\text{Ad}, \text{Tr})$ :

$$\text{Ad}: R\text{-}f\mathcal{B}im_{N-1}^{\text{at}q} \rightarrow R\text{-}f\mathcal{B}im_N^{\text{at}q}$$

$$\text{Ad} \left( \begin{array}{|c|} \hline \text{c} \\ \hline \end{array} \right) =$$


**Idea: Flank them!**


 & should be thought as
 

and things get stuck, e.g.

topologically stuck: 
 & algebraically stuck: 

Skein re



**Partial Hochschild homology (à la Hogancamp ~2015).**  $R\text{-}f\mathcal{B}im_N^{\text{atq}}$  category of (bicomplexes) of  $\mathfrak{q}$ -graded, free  $R_N$ -bimodules. Adjoint pair  $(\text{Ad}, \text{Tr})$ :

$$\text{Ad}: R\text{-}f\mathcal{B}im_{N-1}^{\text{atq}} \rightarrow R\text{-}f\mathcal{B}im_N^{\text{atq}}$$


$$B \mapsto B \otimes_{R_{N-1}} (R_N^e / (x_N \otimes 1 - 1 \otimes x_N)) \quad \longleftrightarrow \quad \text{Ad} \left( \begin{array}{|c|} \hline c \\ \hline \end{array} \right) =$$

**Theorem (after normalization and flanking).**

We get a triply-graded invariant  $\text{HHH}_M^*(\ell) \in \mathbb{k}\text{-Vect}^{\text{atq}}$  for  $\ell \in \mathcal{B}r(g, n)$ , which respects Markov conjugation and stabilization, i.e.

$$\text{HHH}_M^* \left( \begin{array}{|c|} \hline \dots \\ \hline \ell \\ \hline \dots \\ \hline \end{array} \right) \cong \text{HHH}_M^* \left( \begin{array}{|c|} \hline \dots \\ \hline \ell \\ \hline \dots \\ \hline \end{array} \right)$$

$$\text{HHH}_M^* \left( \begin{array}{|c|} \hline e \\ \hline \ell \\ \hline \end{array} \right) \cong \text{HHH}_M^* \left( \begin{array}{|c|} \hline e \\ \hline \ell \\ \hline \end{array} \right) \cong \text{HHH}_M^* \left( \begin{array}{|c|} \hline e \\ \hline \ell \\ \hline \end{array} \right)$$




David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2018, 12/16

The Markov moves on  $\text{Alk}(p, n)$  are conjugation and stabilization.

**Conjugation.**



**Stabilization.**



They are weaker than the Markov moves.


David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2018, 12/16

$\text{con}(r/d)$  twice on a link



Currently known (to the best of my knowledge).		
Genus	type A	type C
$g = 0$	$\text{Alk}(n) \cong \text{AT}(A_{n-1})$	
$g = 1$	$\text{Alk}(1, n) \cong \text{AT}(A_{n-1})$	$\text{Alk}(1, n) \cong \text{AT}(C_{n-1})$
$g = 2$	$\text{Alk}(2, n) \cong \text{AT}(C_{n-1})$	
$g \geq 2$		
And some $\mathbb{Z}/2\mathbb{Z}$ -orbifolds $(\mathbb{Z}/2\mathbb{Z} \curvearrowright \mathbb{C}^2)$ -pairs		
Genus	type D	type B
$g = 0$		
$g = 1$	$\text{Alk}(1, n)_{\text{orb}} \cong \text{AT}(D_{n-1})$	$\text{Alk}(1, n)_{\text{orb}} \cong \text{AT}(B_{n-1})$
$g = 2$	$\text{Alk}(2, n)_{\text{orb}} \cong \text{AT}(D_{n-1})$	$\text{Alk}(2, n)_{\text{orb}} \cong \text{AT}(B_{n-1})$
$g \geq 2$		

(For orbifolds "genus" is just an analogy.)


David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2018, 12/16





David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2018, 12/16



David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2018, 12/16




David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2018, 12/16



David Tubbenhauer, HOMFLYPT homology for links in handlebodies, January 2018, 12/16

There is still much to do...



The Markov moves on  $\mathcal{B}(p, n)$  are conjugation and stabilization.

**Conjugation.**

$$\beta = \alpha R^{-1} \iff \text{for } \epsilon \in \mathcal{B}(p, n), \epsilon \in \{\beta_1, \dots, \beta_{n-1}\}$$

**Stabilization.**


$$\begin{aligned} & \langle \epsilon \cup \{n, n+1\} \rangle \\ \iff & \langle \epsilon \cup \{n\} \cup \{n+1, n+2\} \rangle \iff \end{aligned}$$

They are weaker than the Markov moves.

$\text{con}(\epsilon/\delta)$  twice on a link

Currently known (to the best of my knowledge).		
Genus	type A	type C
$g=0$	$\mathcal{B}(1, n) \cong \text{AT}(A_{n-1})$	
$g=1$	$\mathcal{B}(1, n) \cong \text{AT}(A_{n-1})$	$\mathcal{B}(1, n) \cong \text{AT}(C_{n-1})$
$g=2$	$\mathcal{B}(2, n) \cong \text{AT}(A_{n-1})$	$\mathcal{B}(2, n) \cong \text{AT}(C_{n-1})$
$g \geq 2$		
And some $\mathbb{Z}/2\mathbb{Z}$ -orbifolds $(\mathbb{Z}/2\mathbb{Z} \curvearrowright \mathbb{C})$ -pointures		
Genus	type D	type B
$g=0$		
$g=1$	$\mathcal{B}(1, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \text{AT}(D_{n-1})$	$\mathcal{B}(1, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \text{AT}(B_{n-1})$
$g=2$	$\mathcal{B}(2, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \text{AT}(D_{n-1})$	$\mathcal{B}(2, n)_{\mathbb{Z}/2\mathbb{Z}} \cong \text{AT}(B_{n-1})$
$g \geq 2$		

(For orbifolds "genus" is just an analogy.)



The Markov moves on  $\mathcal{B}(p, n)$  are conjugation and stabilization.

**Theorem (Hiring-Olsberg-Lambropoulos – 2012).**  
Two links in  $\mathcal{B}(p, n)$  are equivalent if and only if they are equal in  $\mathcal{B}(p, n)$  up to conjugation and stabilization.

**Examples.**

for  $p \in \mathbb{C}$


Stabilization:  $\langle \epsilon \cup \{n, n+1\} \rangle \iff \langle \epsilon \cup \{n\} \cup \{n+1, n+2\} \rangle$  for  $\epsilon \in \mathcal{B}(p, n)$

They are weaker than the Markov moves.

**Philosophy 1: Result.** Note that the type A embedding guarantees that any usual invariant of braids produces an invariant of braids in  $\mathcal{B}(p, n)$ .

same resolution:  $\mathbb{N} \times \mathbb{N}$ ,  $\mathbb{N} \times \mathbb{N}$ ,  $\mathbb{N} \times \mathbb{N}$ ,  $\mathbb{N} \times \mathbb{N}$

works quite well (e.g. use Niemeier-Vogt's ideas on the categorified level) but this is for  $M$  being 'large, but finite'



The Coxeter graphs of finite type:

**Examples.**

- Type  $A_n$  → the given a generator-minimal presentation
- Type  $B_n$  → the standard motion preserves the angle between hyperplanes
- Type  $H_1$  → dodecahedron (icosahedron) → exceptional Coxeter group

For  $h_n$  we have a 1-gen:

**Construction:**

- Fix a hyperplane  $H_0$  permitting the adjacent  $i$ -cells of  $\mathcal{F}$
- Fix a hyperplane  $H_1$  permitting two adjacent  $i$ -cells of  $\mathcal{F}$  etc.
- Write a vertex  $v_i$  for each  $H_i$
- Connect  $v_i, v_j$  by an  $n$ -edge for  $H_i, H_j$  having angle  $\pi/n$

**Partial Hochschild homology** (a la Hogeancan – 2015):  $H_1 \mathcal{B}(p, n)_{\text{Hoch}}$  category of  $(\mathbb{Z}/2\mathbb{Z} \curvearrowright \mathbb{C})$  is graded, from  $\mathbb{N}_0$ -knotoids. Adjoint pair  $(\text{Ad}, \text{Tr})$ :

$$\text{Ad}: \mathcal{B}(p, n)_{\text{Hoch}}^{\text{gen}} \rightarrow \mathcal{B}(p, n)_{\text{Hoch}}^{\text{gen}}, \quad \text{Ad} \left( \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$$

**Theorem (after normalization and finiteness).**

We get a triply-graded invariant  $\text{HHH}_1(p) \in \mathbb{Z}\text{-Vec}^{\text{gen}}$  for  $\mathcal{B}(p, n)$ , which respects Markov conjugation and stabilization, i.e.

Thanks for your attention!

The Reidemeister braid relations:



These hold for usual strands only since core strands do not cross each other, e.g.



[← Back](#)


**Brunn  $\sim 1897$ , Alexander  $\sim 1923$ .** For any link  $\ell$  in the 3-ball  $\mathcal{D}^3$  there is a braid in  $\mathcal{B}r(\infty)$  whose closure is isotopic to  $\ell$ .

---

There are various proofs of this result, are all based on the same idea: “Eliminate one by one the arcs of the diagram that have the wrong sense.”

---


Here is an example which works in the context of general 3-manifolds: “Mark the local maxima and minima of the link diagram with respect to some height function and cut open wrong subarcs.”, e.g.




**Markov  $\sim$  1936.** Two links in the 3-ball  $\mathcal{D}^3$  are equivalent if and only if they are equal in  $\mathcal{B}r(\infty)$  up to conjugation and stabilization.

---

### Conjugation.




### Stabilization.




**Markov  $\sim$  1936.** Two links in the 3-ball  $\mathcal{D}^3$  are equivalent if and only if they are equal in  $\mathcal{B}r(\infty)$  up to conjugation and stabilization.


---

### Conjugation.




### Stabilization.





**Figure:** The first ever “published” braid diagram. (Page 283 from Gauß’ handwritten notes, volume seven,  $\leq 1830$ ).

**Tits  $\sim 1961++$ .** Gauß’ braid group is the type A case of more general groups.




Artin's approach: "Arithmetrization of braids".  
 However, he still needs topological arguments.

And this is one main problem why general Artin–Tits groups are so complicated:  
 Basically, they are "infinite groups without extra structure".

Ad. Gauß  
 cd. ob.  
 Man findet mir in jeder mehr je jähre wie oft + mit = verhält

**Figure:** The first ever "published" braid diagram. (Page 283 from Gauß' handwritten notes, volume seven, ≤1830).

**Tits** ~1961++. Gauß' braid group is the type A case of more general groups.



**Figure:** The Coxeter graphs of finite type. (Picture from [https://en.wikipedia.org/wiki/Coxeter\\_group](https://en.wikipedia.org/wiki/Coxeter_group).)

### Examples.


Type  $A_3 \iff$  tetrahedron  $\iff$  symmetric group  $S_4$ .


Type  $B_3 \iff$  cube/octahedron  $\iff$  Weyl group  $(\mathbb{Z}/2\mathbb{Z})^3 \times S_3$ .

Type  $H_3 \iff$  dodecahedron/icosahedron  $\iff$  exceptional Coxeter group.

For  $I_8$  we have a 4-gon:

**Idea (Coxeter ~1934++).**





**Figure:** The Coxeter graphs of finite type. (Picture from [https://en.wikipedia.org/wiki/Coxeter\\_group](https://en.wikipedia.org/wiki/Coxeter_group).)

## Examples.

Type  $A_3 \iff$  tetrahedron  $\iff$  symmetric group  $C_4$


Type  $B_3 \iff$  cube/octaneon  $\iff$  wreath group  $(\mathbb{Z}/2\mathbb{Z}) \ltimes S_3$ .


Type  $H_3 \iff$  dodecahedron/icosahedron  $\iff$  exceptional Coxeter group.

For  $I_8$  we have a 4-gon:

Fix a flag  $F$ .

Idea (Coxeter  $\sim 1934++$ ).





**Figure:** The Coxeter graphs of finite type. (Picture from [https://en.wikipedia.org/wiki/Coxeter\\_group](https://en.wikipedia.org/wiki/Coxeter_group).)

### Examples.

Type  $A_3 \iff$  tetrahedron  $\iff$  symmetric group  $S_4$ .

Type  $B_3 \iff$  cube/octahedron  $\iff$  Weyl group  $(\mathbb{Z}/2\mathbb{Z})^3 \times S_3$ .


Type  $H_3 \iff$  dodecahedron/icosahedron  $\iff$  exceptional Coxeter group.


For  $I_8$  we have a 4-gon:

Fix a flag  $F$ .

Idea (Coxeter  $\sim 1934++$ ).

Fix a hyperplane  $H_0$  permuting the adjacent 0-cells of  $F$ .





**Figure:** The Coxeter graphs of finite type. (Picture from [https://en.wikipedia.org/wiki/Coxeter\\_group](https://en.wikipedia.org/wiki/Coxeter_group).)

## Examples.

Type  $A_3 \iff$  tetrahedron  $\iff$  symmetric group  $S_4$ .

Type  $B_3 \iff$  cube/octahedron  $\iff$  Weyl group  $(\mathbb{Z}/2\mathbb{Z})^3 \times S_3$ .

Type  $H_3 \iff$  dodecahedron/icosahedron  $\iff$  exceptional Coxeter group.


For  $I_8$  we have a 4-gon:


Fix a flag  $F$ .

Idea (Coxeter  $\sim 1934++$ ).

Fix a hyperplane  $H_0$  permuting the adjacent 0-cells of  $F$ .

Fix a hyperplane  $H_1$  permuting the adjacent 1-cells of  $F$ , etc.





**Figure:** The Coxeter graphs of finite type. (Picture from [https://en.wikipedia.org/wiki/Coxeter\\_group](https://en.wikipedia.org/wiki/Coxeter_group).)

## Examples.

Type  $A_3 \iff$  tetrahedron  $\iff$  symmetric group  $S_4$ .

Type  $B_3 \iff$  cube/octahedron  $\iff$  Weyl group  $(\mathbb{Z}/2\mathbb{Z})^3 \times S_3$ .

Type  $H_3 \iff$  dodecahedron/icosahedron  $\iff$  exceptional Coxeter group.

For  $I_8$  we have a 4-gon:


Fix a flag  $F$ .


Idea (Coxeter  $\sim 1934++$ ).

Fix a hyperplane  $H_0$  permuting the adjacent 0-cells of  $F$ .

Fix a hyperplane  $H_1$  permuting the adjacent 1-cells of  $F$ , etc.

Write a vertex  $i$  for each  $H_i$ .





**Figure:** The Coxeter graphs of finite type. (Picture from [https://en.wikipedia.org/wiki/Coxeter\\_group](https://en.wikipedia.org/wiki/Coxeter_group).)

### Examples.

This gives a generator-relation presentation.

Type  $A_3 \leftrightarrow$  tetrahedron  $\leftrightarrow$  symmetric group  $S_4$ .

Type  $B_3 \leftrightarrow$  And the braid relation measures the angle between hyperplanes.

Type  $H_3 \leftrightarrow$  dodecahedron/icosahedron  $\leftrightarrow$  exceptional Coxeter group.

For  $I_8$  we have a 4-gon:

Fix a flag  $F$ .


Idea (Coxeter  $\sim 1934++$ ).

Fix a hyperplane  $H_0$  permuting the adjacent 0-cells of  $F$ .

Fix a hyperplane  $H_1$  permuting the adjacent 1-cells of  $F$ , etc.

Write a vertex  $i$  for each  $H_i$ .

Connect  $i, j$  by an  $n$ -edge for  $H_i, H_j$  having angle  $\cos(\pi/n)$ .



Three gradings:

$\mathfrak{q} \leftrightarrow$  internal

&

$\mathfrak{t} \leftrightarrow$  homological

&

$\mathfrak{a} \leftrightarrow$  Hochschild

---

**Example.** To compute Hochschild cohomology take the Koszul resolution

$$\bigotimes_{i=1}^N \left( R^e = R \otimes R^{\text{op}} \xrightarrow{\cdot(x_i \otimes 1 - 1 \otimes x_i)} \mathfrak{a} \mathfrak{q}^2 R^e \right),$$

Tensor it with  $B$ , gives a complex with differentials  $x_i \otimes 1 - 1 \otimes x_i$ , of which we think as identifying the variables. This gives a chain complex having non-trivial chain groups in  $\mathfrak{a}$ -degree  $0, \dots, n$ . Here the  $i^{\text{th}}$  chain group consists of  $\binom{n}{i}$  copies of  $B$ , with differentials given by the various ways of identifying  $i$  variables. The  $a^{\text{th}}$  cohomology =  $a^{\text{th}}$  Hochschild cohomology.

---

**Example.** If  $B$  is already a  $\mathfrak{t}$ -graded complex, then one can take homology of it and gets “triple  $H$ ”.